

Now Available!



A broad spectrum disinfectant in an easy to use tablet

Easy to Use - Perfect for areas in animal facilities requiring the convenience of a pre-measured, error free disinfecting solution.

Powerful - effective against viruses, gram positive and gram negative bacteria, fungi (molds and yeasts) and mycoplasma.

Flexible Control - inanimate surface, equipment, and fogging (wet misting operations).

- Especially suitable for low volume applications
- Readily soluble effervescent tablet
- Mixed solution is stable for 7 days
- Convenient to store, easy to handle
- Perfume free
- Cost Effective

pharmacal
RESEARCH LABORATORIES, INC.



EFFECTIVE AGAINST THE FOLLOWING PATHOGENS:

BACTERIA

- | | | |
|---------------------------------|---------------------------------|----------------------------------|
| Actinobacillus pleuropneumonia | * Helicobacter pylori | Salmonella enterica |
| Bacillus cereus | * Klebsiella pneumoniae | Salmonella typhimurium |
| * Bordetella avium | * Listeria monocytogenes | Shigella sonnei |
| * Bordetella bronchiseptica | * Moraxella bovis (Pink Eye) | Staphylococcus aureus |
| Brucella abortus | Mycoplasma hypopneumonia | Staphylococcus epidermidis |
| Clostridium perfringens | * Mycoplasma gallisepticum | * Streptococcus equi (Strangles) |
| Campylobacter jejuni | * Mycoplasma mycoides | Streptococcus pyogenes |
| Dermatophilus congolensis | Pasteurella multocida | Streptococcus suis |
| Escherichia coli | Pseudomonas aeruginosa | * Taylorella equigenitalis |
| * Fistulous withers (Poll Evil) | * Pseudomonas mallei (Glanders) | * Treponema hyodysenteriae |
| * Haemophilus somnus | * Pseudomonas vulgaris | |

VIRUSES

- | | | |
|--------------------------------------|--|---|
| * Adenovirus Pneumonia | * Equine Infectious Anemia Virus (Swamp Fever) | * Infectious Salmon Anaemia Virus |
| * African Horse Sickness Virus | * Equine Arteritis Virus | * Infective Bovine Rhinotracheitis Virus |
| * African Swine Fever Virus | Equine Herpes Virus (Type 1) | * Leptospira Canicola Virus |
| Avian Influenza Virus | Herpes Virus Equine (Type 3) | * Maedi-Visna Virus |
| Avian Laryngotracheitis Virus | * Hog Cholera Virus | * Marek's Disease Virus |
| Bovine Adenovirus Type 4 | * Equine Contagious Abortion Virus | * Mouse Parvovirus |
| * Bovine Polyoma Virus | * Equine Papillomatosis Virus | Newcastle Disease Virus |
| * Bovine Pseudocowpox Virus | Equine Influenza Virus (Type A) | * PCV2 Virus (PMWS) |
| * Bovine Viral Diarrhea Virus | * Equine Influenza Virus (The Cough) | * Porcine Parvovirus |
| * Calf Rotavirus | Feline Calicivirus | * Porcine Reproductive and |
| Canine Adenovirus (Canine Hepatitis) | * Feline Herpes Virus | Respiratory Syndrome Virus (PARS) |
| * Canine Coronavirus | * Feline Infectious Peritonitis Virus | * Pseudorabies Virus (Aujeszky's Disease) |
| * Canine Parainfluenza Virus | Feline Panleukopenia Virus | * Rotaviral Diarrhea Virus |
| Canine Parvovirus | * Feline Parvovirus | Simian virus (SV40 Virus) |
| * Chicken Anemia Virus | Feline Rhinotracheitis Virus | * Snakehead rhabdovirus |
| * Coital Exanthema Virus | * Foot and Mouth Disease Virus | * Swine Influenza Virus |
| * Distemper Virus | * Infectious Bronchitis Virus | * Swine Vesicular Disease Virus |
| * Duck Adenovirus | * Infectious Bursal Disease Virus | * Transmissible Gastroenteritis Virus (TGE) |
| * Duck Enteritis Virus | * Infectious Canine Hepatitis Virus | * Turkey Herpes Virus (no hard water) |
| * Egg Drop Syndrome Adenovirus | * Infectious Pancreatic Necrosis Virus | * Turkey Rhinotracheitis Virus |
| | | * Vesicular Stomatitis Virus |

FUNGI

- | | | |
|-------------------------|--------------------------------|----------------------------------|
| * Aspergillus fumigatus | * Microsporium canis | Trichophyton mentagrophytes (2%) |
| * Fusarium moniliforme | * Trichophyton spp. (Ringworm) | * Trichophyton spp. (Mud Fever) |

* Not approved in California for use against these organisms

